



OQBN Health Protocol

Consult with your veterinarian and follow all label directions under a veterinarian-client-patient relationship.

Required Health Program	Option #1* Branding and Weaning	Option #2 Prewearing and Weaning	Option #3 Weaning and Postweaning
Respiratory Viral Vaccination IBR, BVD, BRSV, PI ₃	At Branding Modified-Live Virus (MLV) IBR, BVD, BRSV, PI ₃ At Weaning Modified-Live Virus (MLV) IBR, BVD, BRSV, PI ₃	At 2-6 weeks prior to weaning IBR, BVD, BRSV, PI ₃ At Weaning IBR, BVD, BRSV, PI ₃	At Weaning IBR, BVD, BRSV, PI ₃ 14-28 days later IBR, BVD, BRSV, PI ₃
Pasteurella Pneumonia Vaccination	At Branding <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i> At Weaning <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i>	At 2-6 weeks prior to weaning <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i>	At Weaning <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i>
Clostridial Vaccination	At Branding Clostridial vaccine product. Low-dose (2ml) products are preferred. At Weaning Clostridial vaccine product. Low-dose (2ml) products are preferred	At 2-6 weeks prior to weaning Clostridial vaccine product. Low-dose (2ml) products are preferred At Weaning Clostridial vaccine product. Low-dose (2ml) products are preferred	At Weaning Clostridial vaccine product. Low-dose (2ml) products are preferred 14-28 days later Clostridial vaccine product. Low-dose (2ml) products are preferred

*Kirkpatrick, et al., Journal of American Veterinary Medical Association. 2008; 233: 136-142

Continued on back



Consult with your veterinarian and follow all label directions under a veterinarian-client-patient relationship.

Required Health Program	<u>Option #1</u> Branding and Weaning	<u>Option #2</u> Prewearing and Weaning	<u>Option #3</u> Weaning and Postweaning
Dewormed	Recommended	Recommended	Recommended
Castration	Required and healed	Required and healed	Required and healed
Dehorned	Required and healed	Required and healed	Required and healed
Weaned	At least 45 days	At least 45 days	At least 45 days
Feed Concentrate/ Bunk Training 7 days minimum	Recommended	Recommended	Recommended
Mineral Supplement	Recommended	Recommended	Recommended
Coccidiostat	Recommended	Recommended	Recommended
Verification	21 days before the sale or shipping date	21 days before the sale or shipping date	21 days before the sale or shipping date

Third-party verification is required and can be accomplished by forwarding the enrollment form to Gant Mourer in the OQBN office (see below). Soon after receiving the enrollment form the OQBN representative may conduct a phone audit or a farm visit in the case of first-time participants.

**Oklahoma Quality Beef Network
C/O Gant Mourer
201 Animal Science
Stillwater, OK 74078
405-744-6060
GantM@okstate.edu**