



OQBN Health Protocol

Consult with your veterinarian and follow all label directions under a veterinarian-client-patient relationship.

Required Health Program	Option #1* Branding and Weaning	Option #2 Prewearing and Weaning	Option #3 Weaning and Postweaning
Respiratory Viral Vaccination IBR, BVD, BRSV, PI ₃	At Branding (2-3 months of age) IBR, BVD, BRSV, PI ₃ (modified live virus [MLV])** At Weaning Modified-Live Virus (MLV) IBR, BVD, BRSV, PI ₃	At 2-6 weeks prior to weaning IBR, BVD, BRSV, PI ₃ (MLV)** At Weaning IBR, BVD, BRSV, PI ₃ (MLV)	At Weaning IBR, BVD, BRSV, PI ₃ (MLV) 14-28 days later IBR, BVD, BRSV, PI ₃ (MLV)
Pasteurella Pneumonia Vaccination	At Weaning <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i>	At 2-6 Weeks Prior to Weaning <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i> Or At Weaning <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i>	At Weaning <i>Mannheimia haemolytica</i> or with a combination <i>Mannheimia haemolytica</i> - <i>Pasteurella multocida</i>
Clostridial Vaccination	At Branding Clostridial vaccine product. Low-dose (2ml) products are preferred. At Weaning Clostridial vaccine product. Low-dose (2ml) products are preferred	At 2-6 weeks prior to weaning Clostridial vaccine product. Low-dose (2ml) products are preferred At Weaning Clostridial vaccine product. Low-dose (2ml) products are preferred	At Weaning Clostridial vaccine product. Low-dose (2ml) products are preferred 14-28 days later Clostridial vaccine product. Low-dose (2ml) products are preferred

*Kirkpatrick, et al., Journal of American Veterinary Medical Association. 2008; 233: 136-142

**Consult your veterinarian, use of MLV vaccines on calves pre-weaning is not recommended in herds where cows were not vaccinated with MLV.



Continued on back

Required Health Program	<u>Option #1</u> Branding and Weaning	<u>Option #2</u> Preweaning and Weaning	<u>Option #3</u> Weaning and Postweaning
Dewormed	Recommended	Recommended	Recommended
Castration	Required and healed	Required and healed	Required and healed
Dehorned	Required and healed	Required and healed	Required and healed
Weaned	At least 45 days	At least 45 days	At least 45 days
Feed Concentrate/ Bunk Training 7 days minimum	Recommended	Recommended	Recommended
Mineral Supplement	Recommended	Recommended	Recommended
Coccidiostat	Recommended	Recommended	Recommended
Verification	21 days before the sale or shipping date	21 days before the sale or shipping date	21 days before the sale or shipping date

Consult with your veterinarian and follow all label directions under a veterinarian-client-patient relationship.

Third-party verification is required and can be accomplished by forwarding the enrollment form to the OQBN representative (see following page below). Soon after receiving the enrollment form the OQBN representative may coordinate a phone audit or a farm visit in the case of first-time participants.

Continued on back

OSU Area Livestock Specialists and OQBN Representatives



Northwest OK Area Specialist
 Britt Hicks
 580-349-5439
 britt.hicks@okstate.edu



Northwest OK Area Specialist
 Dana Zook
 580-237-7677
 dana.zook@okstate.edu



Northeast OK Area Specialist
 Earl Ward
 918-686-7800
 earl.ward@okstate.edu



Southwest OK Area Specialist
 Marty New
 580-255-3674
 marty.new@okstate.edu



Southeast OK Area Specialist
 Brian Freking
 580-332-7011
 brian.freking@okstate.edu

